



F1550

FERTILEADER[®] ELITE

9-0-6

FERTILEADER[®] is a foliar fertilizer formulated to support reproductive and late season development, enhance nutrient and water use, and combat abiotic stress.

GUARANTEED ANALYSIS


Total Nitrogen (N)	9%
9% Nitrate Nitrogen	
Soluble Potash (K ₂ O)	6%
Calcium (Ca)	8.5%
Boron (B)	0.1%

DERIVED FROM: Calcium nitrate trihydrate, Potassium hydroxide, Potassium nitrate, Disodium octaborate tetrahydrate.

ALSO CONTAINS BENEFICIAL SUBSTANCES:

4% Seaweed Extracts*

*Derived from *Ascophyllum nodosum*



**PATENTED
LIQUID
FORMULA**

CROPS	APPLICATION STAGE	FOLIAR APPLICATION RATE 1-6 APPLICATIONS
Root & Tuber Vegetables including Potato, Carrot, Onion, Sugarbeet, etc.	During tuber bulking and root crop expansion, prior to row canopy closing	0.5-1.5 qts/ac
Fruits and Fruiting Vegetables including Tomatoes, Peppers, Eggplants, Cucumber, Zucchini, Pumpkins, Squashes, Melon, Strawberry, etc.	At bloom initiation, at fruit development through finishing/size	1-2 qts/ac
Field and Row Crops including Corn, Cotton, Soybean, Wheat, Rice, Peanut, Sorghum, Barley, Canola, Rice, and other Cereal Grains, etc.	At early-to-late reproductive stages starting at flowering through fruit/seed/grain finishing	0.5-1.5 qts/ac
Alfalfa, Clover, Other Grass Crops	Apply 7-14 days prior to harvest, 7-14 days after cutting, at dormancy break	0.5-2 qts/ac
Orchard and Vineyards including but not limited to: Apples, Pears, Peaches, Nectarine, Grapes, Tree Nut, Blueberry, Hops, Cranberry, Citrus	Late bloom to petal fall, at fruit sizing/finishing	1-4 qts/ac
Tobacco, Leafy Crops and Cole Crops (Cabbage, Brussels Sprouts, Broccoli)	During leaf expansion, initiated prior to head formation	1-2 qts/ac

DIRECTIONS FOR USE: Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Shake well before use. Apply only as directed. If foliar applying, dilute into at least 15 gallons/acre. On aerial applications, use minimum 5 gal/ac of water. More frequent applications may be needed to correct deficiencies. A 3-day minimum interval between applications is recommended.

These recommendations are guidelines only. Optimum rate of application will vary depending on environmental conditions, season timing, plant species, and current plant health status. For best results, follow soil and tissue test recommendations.

Always wear appropriate chemical mixing personal protective equipment (PPE), including eye/face and hand protection.

WARNING: This fertilizer carries added Boron and is intended for use only on directed crops. Overapplication or excessive use of Boron, or use on crops other than those listed, can result in Boron toxicity. This may cause foliar damage or serious crop injury. The proper use of soil test recommendations will minimize the risk of overapplication of Boron.

COMPATIBILITY, TANK MIXING AND STORAGE: Check mixture compatibility using a jar test prior to application. Fill mix tank with half of required water and begin agitation. Add products in this order, mixing thoroughly after each addition: adjuvants, pesticides, then fertilizers. Fill tank with remainder of water and continue agitation until solution is completely mixed. Manufacturer responsibility is limited to the guarantee of the product's composition as per label in certified packaging. In all cases, test the compatibility prior to treatment. Store in a cool and dry place. Avoid storing in sunlight or high heat environments. Do not store under 32°F (0°C).

WEIGHT PER GALLON: 12.2 lbs (5.53 kg)

DANGER

H319: Causes serious eye irritation. H335: May cause respiratory irritation. H360: May damage fertility or the unborn child (oral). P202: Do not handle until all safety precautions have been read and understood. P261: Avoid breathing vapors. P280: Wear eye protection, protective clothing, protective gloves. P304+P340: If inhaled: Remove person to fresh air and keep comfortable for breathing. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P313: If exposed or concerned: Get medical advice/attention.

NOTICE OF WARRANTY: TIMAC AGRO USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA, Inc. In no case shall TIMAC AGRO USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risks shall be assumed by the Buyer. TIMAC AGRO USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



PLEASE READ CAREFULLY BEFORE USING
KEEP OUT OF REACH OF CHILDREN
SHAKE WELL BEFORE USE

Manufactured By:
TIMAC AGRO USA Inc.
P.O. Box 888 - Route 724 & I-176
Reading, PA 19607-0888 - USA
us.timacagro.com
800-545-5474

