



TimaCal® is a specialty liquid calcium fertilizer that improves calcium uptake and solves related disorders in fruits, vegetables and grains.

GUARANTEED ANALYSIS

Calcium (Ca) 10.7%

DERIVED FROM: Calcium chloride.

ALSO CONTAINS BENEFICIAL SUBSTANCES:

Seaweed Extracts* 20%

*Derived from *Ascophyllum spp.*

CROPS	APPLICATION STAGE	APPLICATION RATE 1-6 APPLICATIONS
Orchard and Vineyards including but not limited to: Apples, Pears, Peaches, Nectarine, Grapes (Juice, Table, Wine), Citrus, Blueberry, Hops.	Foliar: bud break to harvest	1-3 qts/ac foliar 2-4 qts/ac soil applied as often as bi-weekly
Root and Vegetables including but not limited to: Sugarbeet, Potato, Carrot, Onion, and other root crops, Tomato, Pepper, Eggplant, Cucurbits, Pumpkin, Squashes, Strawberry, Cranberry, Leafy Greens, Cole crops (Cabbage, Brussel Sprouts, Broccoli).	With liquid side-dress or through fertigation (do not mix with sulfur) Foliar: flower to harvest	1-2 qts/ac foliar 2-4 qts/ac soil applied as often as bi-weekly
Field and Row Crops including but not limited to: Corn, Wheat, Barley, Rye, Oats, Sorghum, other cereal grains, Sunflower, Canola, Soybean, Lentils, Dry Beans, Field Peas, Peanuts, Cotton, Rice.	With liquid side-dress (do not mix with sulfur) Foliar: flower to harvest	1-2 qts/ac foliar 2-4 qts/ac soil applied
Alfalfa, Clover, Grass crops.	Foliar apply at spring green up and again 3-7 days after each cutting	1-2 qts/ac foliar 1-4 qts/ac soil applied

These recommendations are guidelines only. Please refer to tissue tests or the advice of your local TIMAC AGRO Field Representative for specific recommendations.

DIRECTIONS FOR USE: Field, Row, Vegetable, Orchard and Vine crops: apply 1-4 quarts per acre per application throughout the growing season. More frequent applications at 1 qt/ac may be needed to correct deficiencies or on crops that have high calcium demand. On aerial applications, use minimum 5 gal/ac of H₂O.

Optimum rate of application will vary depending on soil properties (such as pH, organic matter content, texture), weather conditions, time of year, plant species and general health. For best results, follow soil and tissue test recommendations.

Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Do not apply if a heavy rain is expected. Do not apply near water, storm drains or drainage ditches or to any impervious surfaces. Only apply to the intended application area. Shake well before use.

Always wear appropriate chemical mixing personal protective equipment (PPE), including eye/face and hand protection.

COMPATIBILITY, TANK MIXING AND STORAGE: Check mixture compatibility using a jar test prior to application. Fill mix tank with half of required water and begin agitation. Add products in this order, mixing thoroughly after each addition: adjuvants, pesticides, then fertilizers. Fill tank with remainder of water and continue agitation until solution is completely mixed. Manufacturer responsibility is limited to the guarantee of the product's composition as per label in certified packaging. In all cases, test the compatibility prior to treatment. Store in a cool and dry place, avoid storing in sunlight or high heat environments. Do not store under 32°F (0°C).

WEIGHT PER GALLON: 11.1 lbs (1.33 kg/l)

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN - READ ENTIRE LABEL BEFORE USING THIS PRODUCT - RECIRCULATE BEFORE USE

WARNING

H319: Causes serious eye irritation.

P280: Wear face protection, protective clothing, protective gloves.

P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

P337+P313: If eye irritation persists: Get medical advice/attention.



NOTICE OF WARRANTY: TIMAC AGRO USA INC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA INC. In no case shall TIMAC AGRO USA INC be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risks shall be assumed by the Buyer. TIMAC AGRO USA INC makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Guaranteed By:
TIMAC AGRO USA INC
P.O. Box 888 - Route 724 & I-176
Reading, PA 19607-0888 - USA
800-545-5474

