

MAXIFRUIT



3-7-7

MAXIFRUIT is a foliar fertilizer focused on supporting fruit set, sizing, and quality during the most demanding stages of crop development.

GUARANTEED ANALYSIS

Total Nitrogen (N)	3%	Manganese (Mn)	0.05%
3% Urea Nitrogen (N)		0.05% Chelated Manganese	
Available Phosphate (P ₂ O ₅) ...	7%	Zinc (Zn)	0.1%
Soluble Potash (K ₂ O)	7%	0.1% Chelated Zinc	

DERIVED FROM: Urea, Phosphoric acid, Potassium hydroxide, Manganese EDTA, Zinc EDTA.

CONTAINS BENEFICIAL SUBSTANCES:

Seaweed Extract (derived from *Ascophyllum ssp.*) 10%

CROPS	# OF APP.	APPLICATION STAGE	APPLICATION RATE (QT/AC)
Fruiting Crops	2-3	One week pre-bloom, fruit set	0.5-2 qt/ac
Cereal and Pulse Crops	2-3	One week pre-bloom, fruit set	0.5-1 qt/ac
Vegetables	2-3	One week pre-bloom, fruit set	0.5-2 qt/ac

DIRECTIONS FOR USE: Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Fill tank half full with clean water then add Maxifruit and continue to agitate while mixing. Keep agitation until the end of the application. Use dilution rate in the table above.

These recommendations are guidelines only. Optimum rate of application will vary depending on environmental conditions, season timing, plant species, and current plant health status. For best results, follow soil and tissue test recommendations.

This product is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

Do not apply if a heavy rain is expected. Do not apply near water, storm drains or drainage ditches or to any impervious surfaces. Only apply to the intended application area.

Always wear appropriate chemical mixing personal protective equipment (PPE), including eye/face and hand protection.

COMPATIBILITY AND STORAGE: MAXIFRUIT is compatible with a wide range of agrochemicals except those with mineral oils and products rich in calcium or magnesium. In all cases, test the compatibility prior to treatment. Do not store under 32°F (0°C).

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT THE REACH OF CHILDREN - RECIRCULATE BEFORE USE

DANGER

H314: Causes severe skin burns and eye damage.
H318: Causes serious eye damage.
P260: Do not breathe spray, vapours.
P280: Wear protective clothing, eye protection, face protection, protective gloves.
P301+P330+P331: If swallowed: Rinse mouth. Do NOT induce vomiting.
P303+P361+P353: If on skin (or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower.
P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P310: Immediately call a doctor, a POISON CENTER.



NOTICE OF WARRANTY: TIMAC AGRO USA INC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA INC. In no case shall TIMAC AGRO USA INC be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risks shall be assumed by the Buyer. TIMAC AGRO USA INC makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

WEIGHT PER GALLON: 10.02 lbs (4.54 kg)

Manufactured By:
TIMAC AGRO USA INC
P.O. Box 888 - Route 724 & I-176
Reading, PA 19607-0888 - USA
800-545-5474

