SEACTIV® AXIS

3-18-0

SEACTIV® is a foliar fertilizer formulated to support reproductive and late season development, enhance nutrient and water use, and combat abiotic stress.

GUARANTEED ANALYSIS

Total Nitrogen (N) 3%
1% Nitrate Nitrogen
2% Urea Nitrogen
Available Phosphate (P_2O_5) 18%
Manganese (Mn) 2.5%
Zinc (Zn) 5.7%

DERIVED FROM: Urea, Manganese nitrate, Phosphoric acid, Zinc oxide.

CONTAINS BENEFICIAL SUBSTANCES:
Seaweed Extracts (derived from *Ascophyllum nodosum*) 8%

CROPS	APPLICATION STAGE	FOLIAR APPLICATION RATE 1-6 APPLICATIONS
Root & Tuber Vegetables including Potato, Carrot, Onion, Sugarbeet, etc.	During tuber bulking and root crop expansion, prior to row canopy closing	1-2 qts/ac
Fruits and Fruiting Vegetables including Tomatoes, Peppers, Eggplants, Cucumber, Zucchini, Pumpkins, Squashes, Melon, Strawberry, etc.	At bloom initiation, at fruit development through finishing/size	1-2 qts/ac
Field and Row Crops including Corn, Cotton, Soybean, Wheat, Rice, Peanut, Sorghum, Barley, Canola, Rice, and other Cereal Grains, etc.	At early-to-late reproductive stages starting at flowering through fruit/seed/grain finishing	1-2 qts/ac
Alfalfa, Clover, Other Grass Crops	Apply 7-14 days prior to harvest, 7-14 days after cutting, at dormancy break	1-2 qts/ac
Orchard and Vineyards including but not limited to: Apples, Pears, Peaches, Nectarine, Grapes, Tree Nut, Blueberry, Hops, Cranberry, Citrus	Late bloom to petal fall, at fruit sizing/finishing	1-2 qts/ac
Tobacco, Leafy Crops and Cole Crops (Cabbage, Brussels Sprouts, Broccoli)	During leaf expansion, initiated prior to head formation	1-2 qts/ac

DIRECTIONS FOR USE: Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Shake well before use. Apply only as directed. If foliar applying, dilute into at least 15 gallons/acre. On aerial applications, use minimum 5 gal/ac of water. More frequent applications may be needed to correct deficiencies. A 3-day minimum interval between applications is recommended.

These recommendations are guidelines only. Optimum rate of application will vary depending on environmental conditions, season timing, plant species, and current plant health status. For best results, follow soil and tissue test recommendations.

Do not apply if a heavy rain is expected. Do not apply near water, storm drains or drainage ditches or to any impervious surfaces. Only apply to the intended application area. Always wear appropriate chemical mixing personal protective equipment (PPE), including eye/face and hand protection.

COMPATIBILITY, TANK MIXING AND STORAGE: Check mixture compatibility using a jar test prior to application. Fill mix tank with half of required water and begin agitation. Add products in this order, mixing thoroughly after each addition: adjuvants, pesticides, then fertilizers. Fill tank with remainder of water and continue agitation until solution is completely mixed. Manufacturer responsibility is limited to the guarantee of the product's composition as per label in certified packaging. In all cases, test the compatibility prior to treatment. Store in a cool and dry place. Avoid storing in sunlight or high heat environments. Do not store under 32°F (0°C).

Information regarding the contents and levels of metals in this product is available on the internet at: http://www.aapfco.org/metals.html

WEIGHT PER GALLON: 11.7 lbs (1.4 kg/l)

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN - SHAKE WELL BEFORE USE

DANGER

H314: Causes severe skin burns and eye damage.
H373: May cause damage to organs (brain) through prolonged or repeated exposure (Inhalation).
P260: Do not breathe spray, vapors.
P280: Wear protective clothing, eye and face protection, protective gloves.
P303+P361+P353: If on skin (or hair): Take off immediately all contaminated clothing. Rinse skin with water/shower.
P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P310: Immediately call a doctor, a POISON CENTER.

NOTICE OF WARRANTY: TIMAC AGRO USA INC warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA INC. In no case shall TIMAC AGRO USA INC be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risks shall be assumed by the Buyer. TIMAC AGRO USA INC makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Manufactured By:
TIMAC AGRO USA INC
P.O. Box 888 - Route 724 & I-176
Reading, PA 19607-0888 – USA
800-545-5474

SPECIMEN

Timac AGRO